

# Title

Author  
Institution

## A Block

Define polynomials  $F_n(x)$  by the following:

$$F_n(x) = (-1)^n (1+x^2)^2 e^{\frac{x^2}{2}} \frac{d}{dx} \left( (1+x^2)^{-1} \frac{d^{n-1}}{dx^{n-1}} e^{-\frac{x^2}{2}} \right), \quad n = 1, 2, 3, \dots,$$

The polynomials  $\{F_n(x)\}$  arose when studying an example of potential of the anharmonic oscillator which has an equidistant spectrum.

$$H = -\frac{1}{2} \frac{d^2}{dx^2} + q(x), \quad \text{where } q(x) = \frac{1}{8}x^2 + \frac{2}{1+x^2} - \frac{4}{(1+x^2)^2} + \frac{1}{3}$$

and satisfy the following:

$$F_n(x) = H e_{n+2}(x) + 2(n+2)H e_n(x) + (n+2)(n-1)H e_{n-2}(x)$$

where  $H e_n(x)$  is the  $n$ -th monic Hermite polynomial.

Some math

$$1 + 1 = 2$$

## Another Block

The polynomials  $\{F_n(x)\}$  are orthogonal in the sense that

$$\int_{-\infty}^{\infty} F_n(x) F_m(x) \frac{e^{-\frac{x^2}{2}}}{(1+x^2)^2} dx = (n-1)(n-1)! (2\pi)^{\frac{1}{2}} \delta_{n,m}$$

for any nonnegative integers  $n$  and  $m$ , where  $\delta_{n,m}$  is the Kronecker delta.

The  $\{F_n(x)\}$  are solutions to the following differential equation:

$$(1+x^2) \left( \frac{d^2 F_n(x)}{dx^2} - x \frac{dF_n(x)}{dx} + (n+2)F_n(x) \right) = 4x \frac{dF_n(x)}{dx}$$

## Another Block

**Exceptional Orthogonal Polynomials** are families of polynomials which

- 1 Form a sequence of orthogonal polynomials with finitely many missing degrees.
- 2 Are eigenfunctions of a second order differential operator.
- 3 Are complete in their corresponding weighted  $L^2$  space.

## Example

Here's an example.

## A Block

Let  $\{P_n(x)\}$  be a monic OPS with respect to a symmetric measure  $d\mu(x)$  supported on  $\mathbb{R}$ . We will consider a sequence of polynomials  $R_n(x)$  defined as

$$R_n(x) := P_{n+2}(x) + A_n P_n(x) + B_n P_{n-2}(x), \quad R_0(x) := 1,$$

for sequences  $\{A_n\}$  and  $\{B_n\}$  of real numbers such that

- 1  $R'_n(i) = 0$  for all  $n \geq 1$
- 2  $\int_{-\infty}^{\infty} R_0(x) R_n(x) \frac{d\mu(x)}{(1+x^2)^2} = 0$  for all  $n \geq 1$
- 3  $\int_{-\infty}^{\infty} R_1(x) R_n(x) \frac{d\mu(x)}{(1+x^2)^2} = 0$  for all  $n \geq 2$

Enumerate some math:

- 1  $R_n(x)$  polynomial of degree  $n+2$ .
- 2  $R'_n(-i) = 0$ .
- 3 The sequence  $\{R_n(x)\}$  does not contain a degree 1 or 2 polynomial.

## Alert Block

The polynomials  $R_n(x)$  are orthogonal with respect to  $\frac{d\mu(x)}{(1+x^2)^2}$  i.e.

$$\int_{-\infty}^{\infty} R_n(x) R_m(x) \frac{d\mu(x)}{(1+x^2)^2} = 0$$

for  $n \neq m$  and is nonzero for  $m = n$ .

Let  $\{P_n(x)\}$  be a family of symmetric orthogonal polynomials. Then they satisfy a 3-term recurrence relation which corresponds to the monic Jacobi matrix

$$J = \begin{pmatrix} 0 & 1 & 0 & \dots \\ a_1 & 0 & 1 & \\ 0 & a_2 & 0 & \dots \\ \vdots & & \dots & \dots \end{pmatrix}$$

i.e

$$J\mathbf{P}(x) = x\mathbf{P}(x),$$

where  $\mathbf{P}(x) = (P_0(x), P_1(x), P_2(x), \dots)^\top$ .

## Another Alert Block

Big Theorem

More text.

## Example

Let  $\hat{T}_n(x)$  denote the  $n$ -th monic Chebyshev polynomial.

## Alert Block in an Example Block

There exists sequences of real numbers  $\{A_n\}$  and  $\{B_n\}$  such that the polynomials  $\mathcal{R}_n(x) := \hat{T}_{n+2}(x) + A_n \hat{T}_n(x) + B_n \hat{T}_{n-2}(x)$  satisfy

- 1  $\mathcal{R}'_n(i) = 0$  for all  $n = 1, 2, \dots$
- 2  $\int_{-1}^1 \frac{\mathcal{R}_0(x) \mathcal{R}_n(x)}{\sqrt{1-x^2}(1+x^2)^2} dx = 0$  for all  $n = 1, 2, \dots$
- 3  $\int_{-1}^1 \frac{\mathcal{R}_1(x) \mathcal{R}_n(x)}{\sqrt{1-x^2}(1+x^2)^2} dx = 0$  for all  $n = 2, 3, \dots$

where  $\mathcal{R}_0(x) := 1$ .

Below you can put a picture:




Figure: Caption 1

## References

- 1 V.E. Adler, *On a modification of Crum's method*. Teoret. Mat. Fiz. 101 (1994), no. 3, 323–330 (Russian); translation in Theoret. and Math. Phys. 101 (1994), no. 3, 1381–1386 (1995)
- 2 R. Bailey and M. Derevyagin. *DEK-type orthogonal polynomials and a modification of the Christoffel formula*. J. Comput. Appl. Math. 438 (2024), Paper No. 115561.
- 3 S.Yu. Dubov, V.M. Eleonskii, and N.E. Kulagin, *Equidistant Spectra of Anharmonic Oscillators*, Soviet Phys. JETP 75 (1992), no. 3, 446–451; translated from Zh. Èksper. Teoret. Fiz. 102 (1992), no. 3, 814–825 (Russian).